

Spotrete™ F



Protect Against Rabbit, Deer and Rodent Depredation.

Spotrete™ F protects nursery stock, shrubs, ornamentals, non-bearing orchards and vines such as grape vines from rabbit, deer and rodent depredation. One application will protect treated parts from 1 to 3 months depending upon weather conditions.

KEY PERFORMANCE BENEFITS

- Animal repellent
- Tank-mix compatible with most commonly used pesticides
- Preventive and curative action

EPA REG. NO. 45728-26-1001

ACTIVE INGR. thiram (44%)

FORMULATION flowable

CHEM. FAMILY dithiocarbamate

FRAC NO. M3

REI 24 hours

PACKAGE SIZE 2 x 2.5 gal

APPLICATION spray



FIELD-GROWN NURSERY



CONTAINER-GROWN NURSERY



CAUTION



Nufarm

Grow a better tomorrow.

KEY USES

Animal repellency on deciduous and evergreen trees and shrubs and non-bearing vines and orchards such as grape vines

KEY ANIMALS REPELLED

(See product label for complete list)

Deer
Rabbit
Rodent

USE SUMMARY

DEER, RABBIT AND RODENT REPELLENCY: Spotrete™ F can be used to protect nursery stock, shrubs, ornamentals, non-bearing orchards and vines such as grape vines from rabbit, deer and rodent depredation. Non-bearing is defined as crops that will not bear fruit within one year of application. Spotrete F cannot be used on crops that will bear fruit within one year of application. One application will protect treated parts from 1 to 3 months, depending upon weather conditions.

This product can be used as a repellent in deterring deer from moving into unwanted areas as a result of local feeding and browsing. The product is most effective when used in combination with mechanical deterrents such as ditches, fences, etc. Prepare and spray Spotrete F as indicated below depending upon season and type of application desired. Spray to a height of 5 feet (1.5 meters). Repeat applications may be necessary to protect new growth. The addition of a latex based sticker prolongs the residual effectiveness of the treatment. Repellency is the result of rendering the treated plant parts distasteful. Some feeding may occur.

GROWING SEASON SPRAY APPLICATION: Add 1 quart Spotrete F to 7 quarts of water and apply as a full coverage spray to all portions of the plant to be protected. Spray to runoff. For greater retention and residual efficacy add 1 pint of latex based sticker to the spray solution. Do not use wetting agents or surfactants as foliar damage may result. For best results spray when temperatures are greater than 40°F and relative humidity is less than 70 percent. Coverage depends on plant density.

BRUSH ON APPLICATION: Add Spotrete F undiluted with a paintbrush to stem parts and trunks to be protected. For greater retention and residual efficacy blend 4 fluid ounces of latex based sticker into 1 gallon of Spotrete F prior to brushing.

See label for complete directions for use.

PRECAUTIONS AND RESTRICTIONS

Not for animal repellency use in California.

Spotrete F cannot be used on crops that will bear fruit within one year of application.

The maximum application rate for products applied as a mammal (deer) repellent is one quart of product in 3-7 gallons of water applied in an area of 1000 square feet.

Applications to ornamentals will be restricted to the following 17 northern states and allowed during the winter season only (October thru March): OH, PA, NY, MI, CT, MA, IN, IL, NJ, WV, MN, WI, VT, NH, RI, DE, and MD.

DO NOT graze or feed cover crops from treated orchards.

See label for complete application precautions and restrictions.

